


This listing of claims replaces all prior versions, and listings, of claims in the captioned application.

Listing of Claims:

1. (Original) A compound of formula (I),



the *N*-oxide forms, the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, wherein


n is 0, 1, 2 or 3 and when *n* is 0 then a direct bond is intended;

t is 0, 1, 2, 3 or 4 and when *t* is 0 then a direct bond is intended;

each Q is nitrogen or ;

each X is nitrogen or ;

each Y is nitrogen or ;

each Z is nitrogen or ;

R¹ is -C(O)NR⁷R⁸, -NHC(O)R⁹, -C(O)-C₁₋₆alkanediylSR⁹, -NR¹⁰C(O)N(OH)R⁹, -NR¹⁰C(O)C₁₋₆alkanediylSR⁹, -NR¹⁰C(O)C=N(OH)R⁹ or another Zn-chelating-group

wherein R⁷ and R⁸ are each independently selected from hydrogen, hydroxy, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, aminoC₁₋₆alkyl or aminoaryl;

R⁹ is independently selected from hydrogen, C₁₋₆alkyl, C₁₋₆alkylcarbonyl, arylC₁₋₆alkyl, C₁₋₆alkylpyrazinyl, pyridinone, pyrrolidinone or methylimidazolyl;

R¹⁰ is independently selected from hydrogen or C₁₋₆alkyl;


R² is hydrogen, halo, hydroxy, amino, nitro, C₁₋₆alkyl, C₁₋₆alkyloxy, trifluoromethyl, di(C₁₋₆alkyl)amino, hydroxyamino or naphthalenylsulfonylpyrazinyl;

-L- is a direct bond or a bivalent radical selected from C₁-6alkanediyl,
C₁-6alkanediyoxy, amino, carbonyl or aminocarbonyl;


each R³ independently represents a hydrogen atom and one hydrogen atom can be
replaced by a substituent selected from aryl;

R⁴ is hydrogen, hydroxy, amino, hydroxyc₁-6alkyl, C₁-6alkyl, C₁-6alkyloxy,
arylC₁-6alkyl, aminocarbonyl, hydroxycarbonyl, aminoc₁-6alkyl,
aminocarbonylC₁-6alkyl, hydroxycarbonylC₁-6alkyl, hydroxyaminocarbonyl,
C₁-6alkyloxycarbonyl, C₁-6alkylaminoC₁-6alkyl or di(C₁-6alkyl)aminoC₁-6alkyl;


—(A) is a radical selected from




(a-1)




(a-2)




(a-3)




(a-4)




(a-5)




(a-6)




(a-7)




(a-8)




(a-9)




(a-10)




(a-11)




(a-12)




(a-13)




(a-14)




(a-15)




(a-16)




(a-17)




(a-18)




(a-19)




(a-20)




(a-21)




(a-22)




(a-23)




(a-24)




(a-25)




(a-26)




(a-27)




(a-28)




(a-29)




(a-30)




(a-31)




(a-32)




(a-33)




(a-34)




(a-35)




(a-36)




(a-37)




(a-38)




(a-39)




(a-40)




(a-41)




(a-42)




(a-43)




(a-44)




(a-45)




(a-46)




(a-47)




(a-48)



(a-49)



(a-50)



(a-51)

wherein each s is independently 0, 1, 2, 3, 4 or 5;

each R^5 and R^6 are independently selected from hydrogen; halo; hydroxy; amino; nitro; trihaloC₁₋₆alkyl; trihaloC₁₋₆alkyloxy; C₁₋₆alkyl; C₁₋₆alkyl substituted with aryl and C₃₋₁₀cycloalkyl; C₁₋₆alkyloxy; C₁₋₆alkyloxyC₁₋₆alkyloxy; C₁₋₆alkylcarbonyl; C₁₋₆alkyloxycarbonyl; C₁₋₆alkylsulfonyl; cyanoC₁₋₆alkyl; hydroxyC₁₋₆alkyl; hydroxyC₁₋₆alkyloxy; hydroxyC₁₋₆alkylamino; aminoC₁₋₆alkyloxy; di(C₁₋₆alkyl)aminocarbonyl; di(hydroxyC₁₋₆alkyl)amino; (aryl)(C₁₋₆alkyl)amino; di(C₁₋₆alkyl)aminoC₁₋₆alkyloxy; di(C₁₋₆alkyl)aminoC₁₋₆alkylamino; di(C₁₋₆alkyl)aminoC₁₋₆alkylaminoC₁₋₆alkyl; arylsulfonyl; arylsulfonylamino; aryloxy; aryloxyC₁₋₆alkyl; arylC₂₋₆alkenediyl; di(C₁₋₆alkyl)amino; di(C₁₋₆alkyl)aminoC₁₋₆alkyl; di(C₁₋₆alkyl)amino(C₁₋₆alkyl)amino; di(C₁₋₆alkyl)amino(C₁₋₆alkyl)aminoC₁₋₆alkyl; di(C₁₋₆alkyl)aminoC₁₋₆alkyl(C₁₋₆alkyl)amino; di(C₁₋₆alkyl)aminoC₁₋₆alkyl(C₁₋₆alkyl)aminoC₁₋₆alkyl; aminosulfonylamino(C₁₋₆alkyl)amino; aminosulfonylamino(C₁₋₆alkyl)aminoC₁₋₆alkyl; di(C₁₋₆alkyl)aminosulfonylamino(C₁₋₆alkyl)amino; di(C₁₋₆alkyl)aminosulfonylamino(C₁₋₆alkyl)aminoC₁₋₆alkyl; cyano; thiophenyl; thiophenyl substituted with di(C₁₋₆alkyl)aminoC₁₋₆alkyl(C₁₋₆alkyl)aminoC₁₋₆alkyl, di(C₁₋₆alkyl)aminoC₁₋₆alkyl, C₁₋₆alkylpiperazinylC₁₋₆alkyl,



hydroxyC₁₋₆alkylpiperazinylC₁₋₆alkyl,
hydroxyC₁₋₆alkyloxyC₁₋₆alkylpiperazinylC₁₋₆alkyl,
di(C₁₋₆alkyl)aminosulfonylpiperazinylC₁₋₆alkyl,
C₁₋₆alkyloxypiperidinyl, C₁₋₆alkyloxypiperidinylC₁₋₆alkyl, morpholinylC₁₋₆alkyl,
hydroxyC₁₋₆alkyl(C₁₋₆alkyl)aminoC₁₋₆alkyl, or di(hydroxyC₁₋₆alkyl)aminoC₁₋₆alkyl;
furanyl; furanyl substituted with hydroxyC₁₋₆alkyl; benzofuranyl; imidazolyl;
oxazolyl; oxazolyl substituted with aryl and C₁₋₆alkyl; C₁₋₆alkyltriazolyl; tetrazolyl;
pyrrolidinyl; pyrrolyl; piperidinylC₁₋₆alkyloxy; morpholinyl; C₁₋₆alkylmorpholinyl;
morpholinylC₁₋₆alkyloxy;
morpholinylC₁₋₆alkyl; morpholinylC₁₋₆alkylamino;
morpholinylC₁₋₆alkylaminoC₁₋₆alkyl; piperazinyl; C₁₋₆alkylpiperazinyl;
C₁₋₆alkylpiperazinylC₁₋₆alkyloxy; piperazinylC₁₋₆alkyl;
naphtalenylsulfonylpiperazinyl; naphtalenylsulfonylpiperidinyl; naphtalenylsulfonyl;
C₁₋₆alkylpiperazinylC₁₋₆alkyl; C₁₋₆alkylpiperazinylC₁₋₆alkylamino;
C₁₋₆alkylpiperazinylC₁₋₆alkylaminoC₁₋₆alkyl; C₁₋₆alkylpiperazinylsulfonyl;
aminosulfonylpiperazinylC₁₋₆alkyloxy; aminosulfonylpiperazinyl;
aminosulfonylpiperazinylC₁₋₆alkyl; di(C₁₋₆alkyl)aminosulfonylpiperazinyl;
di(C₁₋₆alkyl)aminosulfonylpiperazinylC₁₋₆alkyl; hydroxyC₁₋₆alkylpiperazinyl;
hydroxyC₁₋₆alkylpiperazinylC₁₋₆alkyl; C₁₋₆alkyloxypiperidinyl;
C₁₋₆alkyloxypiperidinylC₁₋₆alkyl; piperidinylaminoC₁₋₆alkylamino;
piperidinylaminoC₁₋₆alkylaminoC₁₋₆alkyl;
(C₁₋₆alkylpiperidinyl)(hydroxyC₁₋₆alkyl)aminoC₁₋₆alkylamino;
(C₁₋₆alkylpiperidinyl)(hydroxyC₁₋₆alkyl)aminoC₁₋₆alkylaminoC₁₋₆alkyl;
hydroxyC₁₋₆alkyloxyC₁₋₆alkylpiperazinyl;
hydroxyC₁₋₆alkyloxyC₁₋₆alkylpiperazinylC₁₋₆alkyl;
(hydroxyC₁₋₆alkyl)(C₁₋₆alkyl)amino; (hydroxyC₁₋₆alkyl)(C₁₋₆alkyl)aminoC₁₋₆alkyl;
hydroxyC₁₋₆alkylaminoC₁₋₆alkyl; di(hydroxyC₁₋₆alkyl)aminoC₁₋₆alkyl;
pyrrolidinylC₁₋₆alkyl; pyrrolidinylC₁₋₆alkyloxy; pyrazolyl; thiopyrazolyl; pyrazolyl
substituted with two substituents selected from C₁₋₆alkyl or trihaloC₁₋₆alkyl;
pyridinyl; pyridinyl substituted with C₁₋₆alkyloxy, aryloxy or aryl; pyrimidinyl;
tetrahydropyrimidinylpiperazinyl; tetrahydropyrimidinylpiperazinylC₁₋₆alkyl;
quinolinyl; indole; phenyl; phenyl substituted with one, two or three substituents
independently selected from halo, amino, nitro, C₁₋₆alkyl, C₁₋₆alkyloxy,
hydroxyC₁₋₄alkyl, trifluoromethyl, trifluoromethyloxy, hydroxyC₁₋₄alkyloxy,
C₁₋₄alkylsulfonyl, C₁₋₄alkyloxyC₁₋₄alkyloxy, C₁₋₄alkyloxycarbonyl,
aminoC₁₋₄alkyloxy, di(C₁₋₄alkyl)aminoC₁₋₄alkyloxy, di(C₁₋₄alkyl)amino,
di(C₁₋₄alkyl)aminocarbonyl, di(C₁₋₄alkyl)aminoC₁₋₄alkyl,






di(C₁₋₄alkyl)aminoC₁₋₄alkylaminoC₁₋₄alkyl,
di(C₁₋₄alkyl)amino(C₁₋₄alkyl)amino, di(C₁₋₄alkyl)amino(C₁₋₄alkyl)aminoC₁₋₄alkyl,
di(C₁₋₄alkyl)aminoC₁₋₄alkyl(C₁₋₄alkyl)amino,
di(C₁₋₄alkyl)aminoC₁₋₄alkyl(C₁₋₄alkyl)aminoC₁₋₄alkyl,
aminosulfonylamino(C₁₋₄alkyl)amino,
aminosulfonylamino(C₁₋₄alkyl)aminoC₁₋₄alkyl,
di(C₁₋₄alkyl)aminosulfonylamino(C₁₋₄alkyl)amino,
di(C₁₋₄alkyl)aminosulfonylamino(C₁₋₄alkyl)aminoC₁₋₆alkyl, cyano,
piperidinylC₁₋₄alkyloxy, pyrrolidinylC₁₋₄alkyloxy, aminosulfonylpiperazinyl,
aminosulfonylpiperazinylC₁₋₄alkyl, di(C₁₋₄alkyl)aminosulfonylpiperazinyl,
di(C₁₋₄alkyl)aminosulfonylpiperazinylC₁₋₄alkyl, hydroxyC₁₋₄alkylpiperazinyl,
hydroxyC₁₋₄alkylpiperazinylC₁₋₄alkyl, C₁₋₄alkyloxypiperidinyl,
C₁₋₄alkyloxypiperidinylC₁₋₄alkyl, hydroxyC₁₋₄alkyloxyC₁₋₄alkylpiperazinyl,
hydroxyC₁₋₄alkyloxyC₁₋₄alkylpiperazinylC₁₋₄alkyl,
(hydroxyC₁₋₄alkyl)(C₁₋₄alkyl)amino, (hydroxyC₁₋₄alkyl)(C₁₋₄alkyl)aminoC₁₋₄alkyl,
di(hydroxyC₁₋₄alkyl)amino, di(hydroxyC₁₋₄alkyl)aminoC₁₋₄alkyl, furanyl,
furanyl substituted with -CH=CH-CH=CH-, pyrrolidinylC₁₋₄alkyl, pyrrolidinylC₁₋₄alkyloxy,
morpholinyl, morpholinylC₁₋₄alkyloxy, morpholinylC₁₋₄alkyl,
morpholinylC₁₋₄alkylamino, morpholinylC₁₋₄alkylaminoC₁₋₄alkyl, piperazinyl,
C₁₋₄alkylpiperazinyl, C₁₋₄alkylpiperazinylC₁₋₄alkyloxy, piperazinylC₁₋₄alkyl,
C₁₋₄alkylpiperazinylC₁₋₄alkyl, C₁₋₄alkylpiperazinylC₁₋₄alkylamino,
C₁₋₄alkylpiperazinylC₁₋₄alkylaminoC₁₋₆alkyl, tetrahydropyrimidinylpiperazinyl,
tetrahydropyrimidinylpiperazinylC₁₋₄alkyl, piperidinylaminoC₁₋₄alkylamino,
piperidinylaminoC₁₋₄alkylaminoC₁₋₄alkyl,
(C₁₋₄alkylpiperidinyl)(hydroxyC₁₋₄alkyl)aminoC₁₋₄alkylamino,
(C₁₋₄alkylpiperidinyl)(hydroxyC₁₋₄alkyl)aminoC₁₋₄alkylaminoC₁₋₄alkyl,
pyridinylC₁₋₄alkyloxy,
hydroxyC₁₋₄alkylamino, hydroxyC₁₋₄alkylaminoC₁₋₄alkyl,
di(C₁₋₄alkyl)aminoC₁₋₄alkylamino, aminothiadiazolyl,
aminosulfonylpiperazinylC₁₋₄alkyloxy, or thiophenylC₁₋₄alkylamino;
each R⁵ and R⁶ can be placed on the nitrogen in replacement of the hydrogen;

aryl in the above is phenyl, or phenyl substituted with one or more substituents each independently selected from halo, C₁₋₆alkyl, C₁₋₆alkyloxy, trifluoromethyl, cyano or hydroxycarbonyl.


2. (Original) A compound as claimed in claim 1 wherein n is 1 or 2; t is 0, 1, 2 or 4; each Q is $\text{---C}\equiv$; R^1 is ---C(O)NH(OH) ; R^2 is hydrogen or nitro; -L- is a direct bond or a bivalent radical selected from C_{1-6} alkanediyl; R^4 is hydrogen; $\text{---}\text{A}$ is a radical selected from (a-1), (a-2), (a-3), (a-5), (a-6), (a-11), (a-18), (a-20), (a-21), (a-32), (a-33), (a-47) or (a-51); each s is independently 0, 1, 2, or 4; each R^5 and R^6 are independently selected from hydrogen; halo; trihalo C_{1-6} alkyl; C_{1-6} alkyl; C_{1-6} alkyl substituted with aryl and C_{3-10} cycloalkyl; C_{1-6} alkyloxy; C_{1-6} alkylcarbonyl; benzofuranyl; naphthalenylsulfonyl; pyridinyl substituted with aryloxy; phenyl; or phenyl substituted with one substituent independently selected from hydroxy C_{1-4} alkyl or morpholinyl C_{1-4} alkyl.
3. (Original) A compound as claimed in claim 1 wherein t is 1, 2, 3, or 4; R^1 is $\text{---C(O)NR}^7\text{R}^8$, ---C(O)---C_{1-6} alkanediyl SR^9 , $\text{---NR}^{10}\text{C(O)N(OH)R}^9$, $\text{---NR}^{10}\text{C(O)C}_{1-6}$ alkanediyl SR^9 , $\text{---NR}^{10}\text{C(O)C=N(OH)R}^9$ or another Zn-chelating-group wherein R^7 and R^8 are each independently selected from hydrogen, hydroxy, hydroxy C_{1-6} alkyl or amino C_{1-6} alkyl; R^2 is hydrogen, halo, hydroxy, amino, nitro, C_{1-6} alkyl, C_{1-6} alkyloxy, trifluoromethyl or di(C_{1-6} alkyl)amino; -L- is a direct bond or a bivalent radical selected from C_{1-6} alkanediyl, C_{1-6} alkanediyl, amino or carbonyl; R^4 is hydrogen, hydroxy, amino, hydroxy C_{1-6} alkyl, C_{1-6} alkyl, C_{1-6} alkyloxy, aryl C_{1-6} alkyl, aminocarbonyl, amino C_{1-6} alkyl, C_{1-6} alkylamino C_{1-6} alkyl or di(C_{1-6} alkyl)amino C_{1-6} alkyl; $\text{---}\text{A}$ is a radical selected from (a-1), (a-3), (a-4), (a-5), (a-6), (a-7), (a-8), (a-9), (a-10), (a-11), (a-12), (a-13), (a-14), (a-15), (a-16), (a-17), (a-18), (a-19), (a-20), (a-21), (a-22), (a-23), (a-24), (a-25), (a-26), (a-28), (a-29), (a-30), (a-31), (a-32), (a-33), (a-34), (a-35), (a-36), (a-37), (a-38), (a-39), (a-40), (a-41), (a-42), (a-44), (a-45), (a-46), (a-47), (a-48) and (a-51); each s is independently 0, 1, 2, 3 or 4; R^5 is hydrogen; halo; hydroxy; amino; nitro; trihalo C_{1-6} alkyl; trihalo C_{1-6} alkyloxy; C_{1-6} alkyl; C_{1-6} alkyloxy; C_{1-6} alkylcarbonyl; C_{1-6} alkyloxycarbonyl; C_{1-6} alkylsulfonyl; hydroxy C_{1-6} alkyl; aryloxy; di(C_{1-6} alkyl)amino; cyano; thiophenyl; furanyl; furanyl substituted with hydroxy C_{1-6} alkyl; benzofuranyl; imidazolyl; oxazolyl; oxazolyl substituted with aryl and C_{1-6} alkyl; C_{1-6} alkyltriazolyl; tetrazolyl; pyrrolidinyl; pyrrolyl; morpholinyl;

- C₁-6alkylmorpholinyl; piperazinyl;
C₁-6alkylpiperazinyl; hydroxyC₁-6alkylpiperazinyl;
C₁-6alkyloxypiperidinyl; pyrazolyl; pyrazolyl substituted with one or two substituents selected from C₁-6alkyl or trihaloC₁-6alkyl; pyridinyl; pyridinyl substituted with C₁-6alkyloxy, aryloxy or aryl; pyrimidinyl; quinolinyl; indole; phenyl; or phenyl substituted with one or two substituents independently selected from halo, C₁-6alkyl, C₁-6alkyloxy or trifluoromethyl;
- R⁶ is hydrogen; halo; hydroxy; amino; nitro; trihaloC₁-6alkyl; trihaloC₁-6alkyloxy; C₁-6alkyl; C₁-6alkyloxy; C₁-6alkylcarbonyl; C₁-6alkyloxycarbonyl; C₁-6alkylsulfonyl; hydroxyC₁-6alkyl; aryloxy; di(C₁-6alkyl)amino; cyano; pyridinyl; phenyl; or phenyl substituted with one or two substituents independently selected from halo, C₁-6alkyl, C₁-6alkyloxy or trifluoromethyl.
4. (Currently Amended) A compound as claimed in claim 1 ~~and 2~~ wherein n is 1; t is 0 or 1; each Q is $\text{—C}\begin{smallmatrix} \diagup \\ \diagdown \end{smallmatrix}$; each X is nitrogen; each Y is nitrogen; R¹ is —C(O)NH(OH) ; R² is hydrogen; -L- is a direct bond; each R³ independently represents a hydrogen atom; R⁴ is hydrogen; $\text{—}\bigcirc\text{(A)}$ is a radical selected from (a-6), (a-11), (a-20), (a-47) or (a-51); each s is independently 0, 1, or 4; and each R⁵ and R⁶ are independently selected from hydrogen; C₁-6alkyl; C₁-6alkyloxy; naphthalenylsulfonyl; or phenyl substituted with hydroxyC₁-4alkyl or morpholinylC₁-4alkyl.
5. (Currently Amended) A compound ~~according to claim 1, 2 and 4~~ selected from the group consisting of compounds No. 3, No. 4, No. 8, No. 5, No. 7, No. 6 and No. 9.

 <p>0.91-C₂HF₃O₂; Co. No. 3</p>	 <p>0.86-C₂HF₃O₂; Co. No. 4</p>
---	--

	
$C_2HF_3O_2$ (1:1); Co. No.8	$0.83 C_2HF_3O_2$; Co. No.5
	
$-0.79 C_2HF_3O_2$; Co. No.7	$0.83 C_2HF_3O_2$; Co. No.6
	
$0.47 H_2O$ $-1.99 C_2HF_3O_2$; Co. No.9	

6. (Currently Amended) A pharmaceutical composition comprising pharmaceutically acceptable carriers and as an active ingredient a therapeutically effective amount of a compound according to as claimed in claim 1 to 5.
7. (Currently Amended) A process of preparing a pharmaceutical composition as claimed in claim 6 wherein the pharmaceutically acceptable carriers and the a compound according to as claimed in claim 1 to 5 are intimately mixed.
8. (Cancelled)
9. (Cancelled)
10. (Original) A process for preparing a compound as claimed in claim 1, characterized by reacting an intermediate of formula (II) with an appropriate acid, such as for example, trifluoro acetic acid, yielding a hydroxamic acid of formula (I-a), wherein R^1 is $-C(O)NH(OH)$



11. (Original) A method of detecting or identifying a HDAC in a biological sample comprising detecting or measuring the formation of a complex between a labelled compound as defined in claim (I) and a HDAC.
12. (Cancelled)